



## Original article

## Extracellular phospholipase activity of *Malassezia* strains isolated from individuals with and without dermatological disease

Gabriella Pini\*, Elisabetta Faggi

Department of Public Health, University of Florence, Florence, Italy

## ARTICLE INFO

## Article history:

Received 31 January 2011

Accepted 3 May 2011

Available online 19 May 2011

## Keywords:

*Malassezia* species

Phospholipase activity

Skin lesions

## ABSTRACT

**Background:** The *Malassezia* genus includes mainly lipophilic yeasts belonging to the cutaneous microbiota of man and other mammals. Some *Malassezia* species have been associated with various dermatological diseases. The factors permitting the transformation of yeasts of the *Malassezia* genus from a commensal organism to a pathogenic agent are still little known but the production of various enzymes such as lipase, phospholipase and lipoxygenase could contribute to the pathogenic activity of these yeasts.

**Aims:** Here we have determined and compared the extracellular phospholipase activity of sixty human isolates of *Malassezia* so as to relate this feature to the species of *Malassezia* and to the origin (from dermatological diseases or not) of the strains examined.

**Methods:** Phospholipase production was determined using the semi-quantitative egg-yolk plate method.

**Results and conclusions:** *Malassezia obtusa*, *Malassezia slooffiae*, *Malassezia globosa*, *Malassezia restricta* had difficulty developing in the chosen culture medium so that it was not possible to measure phospholipase activity. *Malassezia pachydermatis* showed the highest phospholipase activity. Twenty-nine *Malassezia sympodialis* strains produced phospholipase; the isolates from patients with pityriasis versicolor had significantly higher phospholipase activity than those isolated from healthy individuals. This observation suggests that the phospholipase activity of *Malassezia* may play a role in the onset of skin lesions, especially in the case of pityriasis versicolor.

© 2011 Revista Iberoamericana de Micología. Published by Elsevier España, S.L. All rights reserved.

### Actividad fosfolipasa extracelular de aislamientos de *Malassezia* de individuos con y sin dermatosis

## RESUMEN

**Antecedentes:** El género *Malassezia* incluye numerosas levaduras lipofílicas que pertenecen a la microbiota cutánea del hombre y de otros mamíferos. Algunas especies se han asociado con diversas dermatosis. Apenas se conocen los factores que permiten la transformación de las levaduras del género *Malassezia* de un microorganismo comensal a uno patógeno, pero la producción de diversas enzimas, como lipasas, fosfolipasas y lipooxigenasa, podría contribuir a la actividad patogénica de estas levaduras.

**Objetivos:** En la presente investigación hemos determinado y comparado la actividad de la fosfolipasa extracelular a partir de 60 aislamientos de *Malassezia* en seres humanos para relacionar esta característica con las especies del género y con el origen (dermatosis o no) de las cepas examinadas.

**Métodos:** La producción de fosfolipasa se determinó utilizando el método semicuantitativo de la placa de yema de huevo.

**Resultados y conclusiones:** *Malassezia obtusa*, *Malassezia slooffiae*, *Malassezia globosa* y *Malassezia restricta* crecieron mal en el medio de cultivo seleccionado, por lo que no fue posible determinar la actividad fosfolipasa. *Malassezia pachydermatis* mostró la mayor actividad enzimática. Produjeron fosfolipasa 29 cepas de *Malassezia sympodialis*; los aislamientos de pacientes con pitiriasis versicolor se caracterizaron por una actividad fosfolipasa significativamente mayor que los aislamientos de individuos sanos. Esta observación sugiere que la actividad fosfolipasa de *Malassezia* podría desempeñar un papel en el inicio de las lesiones cutáneas, en particular en el caso de la mencionada micosis.

© 2011 Revista Iberoamericana de Micología. Publicado por Elsevier España, S.L. Todos los derechos reservados.

## Palabras clave:

Especies de *Malassezia*

Actividad fosfolipasa

Lesiones cutáneas

\* Corresponding author.

E-mail address: [gpini@unifi.it](mailto:gpini@unifi.it) (G. Pini).

The *Malassezia* genus includes mainly lipophilic yeasts belonging to the cutaneous microbiota of man and other mammals. These yeasts were first described by Eichstedt in 1846 as being associated with lesions of pityriasis versicolor.

The taxonomy and nomenclature of the *Malassezia* genus has, in recent years, been revised several times. In fact, up until 1995 only three species had been identified: *Malassezia furfur*, *Malassezia pachydermatis* and *Malassezia sympodialis*. The species *Malassezia globosa*, *Malassezia restricta*, *Malassezia obtusa*, and *Malassezia slooffiae* were described in 1995 on the basis of their morphology, ultrastructure, physiology and genomic features.<sup>11,13</sup> In the last few years, the new lipid-dependent species *Malassezia dermatis*,<sup>31</sup> *Malassezia japonica*,<sup>30</sup> and *Malassezia yamatoensis*<sup>29</sup> were isolated from human skin. Other species were found on animals: *Malassezia nana*<sup>15</sup> on cats and dogs, *Malassezia equina*, *Malassezia caprae* and *Malassezia cuniculi*,<sup>3,4</sup> found mainly on horses, goats and rabbits respectively.

Yeasts belonging to the *Malassezia* genus have been associated with various dermatological diseases: pityriasis versicolor (PV), dandruff, seborrheic dermatitis (SD), atopic dermatitis, folliculitis, psoriasis, onychomycosis, and blepharitis. The pathogenic role of *Malassezia* in PV has now been universally accepted although authors disagree about which species is most widely associated with the disease. Some of them suggest that *M. globosa* is the causal agent of PV.<sup>8,9,14</sup>

As well as these diseases, cases of fungemia caused by *M. pachydermatis* and *M. furfur* have been reported in premature newborns and immunocompromised patients artificially fed with lipid emulsions.<sup>32</sup>

Most *Malassezia* species are lipid-dependent; as a result these yeasts probably require lipolytic enzymes such as lipases and phospholipases to use the environmental lipids essential for their growth. In a recent work, Xu<sup>35</sup> described the genome and secretory proteome of two species of *Malassezia*, suggesting that the lipid-dependency of these species is associated with the apparent absence of the gene for fatty acid synthase. Further studies have demonstrated extracellular production of phospholipase and lipase.<sup>7,17</sup>

Phospholipases are a heterogeneous group of enzymes which hydrolyse one or more ester bonds of glycerol-phospholipids. Phospholipids and proteins are the main chemical constituents of the cellular membranes of the host so that the phospholipases and proteases are involved in the destruction processes of such cellular membranes. Many fungal species, such as *Candida albicans*, *Cryptococcus neoformans* and *Aspergillus fumigatus* can produce enzymes belonging to the phospholipases group.<sup>2,20,34</sup> Researchers have associated some extracellular phospholipases with the virulence of *C. albicans*.<sup>16,23</sup>

The factors permitting the transformation of yeasts of the *Malassezia* genus from a commensal organism to a pathogenic agent are still little known, but the production of various enzymes such as lipase, phospholipase and lipoxygenase could contribute to the pathogenic activity of these yeasts too.

Little information is available in literature regarding the phospholipase activity of species of the *Malassezia* genus. Using the Price method (egg yolk medium),<sup>26</sup> extracellular phospholipase activity was demonstrated *in vitro* on strains of *M. furfur* isolated from skin lesions in humans.<sup>22,27</sup> These works were prior to the classification introduced by Guého et al. so, bearing in mind the current classification, some strains included in these studies could be reclassified as different species.

The phospholipase activity of *M. pachydermatis* has also been associated with skin lesions in dogs.<sup>5,6</sup>

In this work we have analysed and compared the extracellular phospholipase activity of *Malassezia* isolates belonging to different species, identified according to the current classification so as to verify whether the various species of *Malassezia* have different phospholipase activity and to relate this feature to the origin (from dermatological diseases or not) of the strains examined.

## Materials and methods

The strains included in this study were isolated from patients with PV or SD, both from skin lesions and from healthy skin, and from individuals without any dermatological disease. Fourteen reference strains from the Pasteur Institute (IP) and Central Bureau voor Schimmelcultures (CBS) (Table 1) were also examined.

Our strains were isolated from scaly skin. The samples were inoculated on Dixon modified (m-Dixon) agar,<sup>12</sup> supplemented with 0.05 g chloramphenicol pH 6.0.

The inoculated plates were incubated aerobically at 32 °C for 2 weeks in plastic boxes with a layer of paper soaked in water on the bottom to prevent desiccation. The culture was considered positive when at least one *Malassezia* colony was observed. *Malassezia* yeasts growing on m-Dixon agar were selected and subcultured for species identification. *Malassezia* species were identified by their morphological characteristics, catalase test, and growth in the presence of Tween 20, 40, 60 and 80 as unique lipid supplementation, using the scheme established by Guillot et al.<sup>12</sup> To improve the differential identification of *M. furfur*, *M. sympodialis* and *M. slooffiae*, the splitting of esculin ( $\beta$ -glucosidase activity), the assimilation of Cremophor EL and tryptophan described by Mayser et al. were also assayed.<sup>19,21</sup>

Phospholipase production was determined using the semi-quantitative egg-yolk plate method previously reported.<sup>25,26</sup> The egg-yolk plates were inoculated with 5  $\mu$ l of a fresh culture containing 10<sup>8</sup> cells/ml. The inoculated egg-yolk plates were incubated at 32 °C in plastic boxes as described above, and readings were taken daily from day 7 to day 20. The formation of precipitation zones around the colony was considered indicative of enzyme production.

The production of phospholipase (Pz) was expressed as a ratio of colony diameter to total diameter of the colonies plus the precipitation zone.<sup>25</sup> When the Pz was equal to 1.0 the samples were considered negative, and when the Pz was less than 1.0, phospholipase activity was considered positive. Each strain was tested in duplicate and the Pz value represents an average of the two

**Table 1**  
Number, species and origin of the strains examined.

Species	Number of examined strains	PV patients skin with lesions	PV patients skin without lesions	SD patients skin with lesions	SD patients skin without lesions	Healthy individuals	IP or CBS
<i>M. furfur</i>	5						5
<i>M. pachydermatis</i>	1						1
<i>M. obtusa</i>	1						1
<i>M. slooffiae</i>	4			1			3
<i>M. globosa</i>	17	6		4	1	5	1
<i>M. restricta</i>	1			1			
<i>M. sympodialis</i>	31	4	3	2	4	15	3
Total	60	10	3	8	5	20	14

CBS: Central Bureau voor Schimmelcultures; IP: Pasteur Institute; PV: pityriasis versicolor; SD: seborrheic dermatitis.

**Table 2**Phospholipasic activity of the *Malassezia* species expressed as a Pz value mean after 15 and 20 days of culture.

Species	Strains examined	Strains with positive development	Strains with positive phospholipase	Pz 15 days	Pz 20 days
<i>M. furfur</i>	5	5	1	1	0.68
<i>M. pachydermatis</i>	1	1	1	0.65	0.53
<i>M. obtusa</i>	1	0	0		
<i>M. slooffiae</i>	4	3 <sup>a</sup>	0		
<i>M. globosa</i>	17	8 <sup>a</sup>	0		
<i>M. restricta</i>	1	0	0		
<i>M. sympodialis</i>	31	31	29	0.68	0.62

Pz: ratio of colony diameter to total diameter of the colonies plus the precipitation zone.

<sup>a</sup> Weak development.

Pz values reported. The Student *t* test was used for statistical analysis of the Pz value; a *P* value  $\leq 0.05$  was considered significant.

## Results

The phospholipasic activity determined by the Price method is a simple technique applicable to a large number of strains. However, the measurement of the precipitation zone was not easy; in fact, the colony was already visible after 48 hours of incubation, while the precipitation zone around the colony was not observed until 8–10 days later, appearing extremely dense but clearly distinguishable from the colony. Up until day 15 of incubation both the diameter of the colony and that of the precipitation zone increased, whereas from day 15 to day 20 only the precipitation zone increased. Measurements of the precipitation zone taken at the day 15 and at day 20 from incubation were therefore deemed significant.

The ability of the species belonging to the *Malassezia* genus to produce phospholipase is shown in Table 2.

*M. furfur* grew well on this medium but only for one strain it was possible to distinguish a late (day 20) precipitation zone around the colony. *M. pachydermatis* grew well and showed phospholipasic activity. *M. sympodialis* grew well and showed phospholipasic activity with the exception of two strains, one isolated from healthy subjects, the other supplied by IP, ex human normal skin.

*M. obtusa*, *M. slooffiae*, *M. globosa*, *M. restricta*, showed poor growth and did not produce a precipitate.

The greatest phospholipasic activity (Pz value) was shown by the strain *M. pachydermatis* after 20 days of incubation.

For *M. sympodialis* we compared the phospholipasic activity expressed as a Pz value mean with the dermatological disease presented by the patient.

Table 3 shows the Pz mean value of the strains isolated from patients with PV or SD, from individuals without any dermatological disease and from the reference strains of IP and CBS.

The Student *t* test highlighted significant differences between the Pz values of strains isolated from patients with PV and those of strains isolated from healthy individuals, both for readings on day 15 and day 20 subsequent to incubation ( $P=0.04$  and  $0.05$  respectively), while such difference was not significant for strains isolated from patients with SD ( $P=0.08$  and  $0.4$ ). The CBS and IP reference

**Table 3**Phospholipasic activity of *M. sympodialis* expressed as a Pz mean value after 15 and 20 days of culture.

Strains origin	Pz 15 days	Pz 20 days
PV patients	0.63 <sup>a</sup>	0.58 <sup>a</sup>
SD patients	0.69	0.65
Healthy individuals	0.73	0.66
IP or CBS	0.73	0.67

Pz: ratio of colony diameter to total diameter of the colonies plus the precipitation zone; CBS: Central Bureau voor Schimmelcultures; IP: Pasteur Institute; PV: pityriasis versicolor; SD: seborrheic dermatitis.

<sup>a</sup> Statistically significant differences ( $P \leq .05$ ).**Table 4**Phospholipasic activity of *M. sympodialis* with reference to strains origin.

Strains origin	Pz 15 days	Pz 20 days
PV patients lesioned skin	0.63	0.58
PV patients healthy skin	0.64	0.59
SD patients lesioned skin	0.69	0.60 <sup>a</sup>
SD patients healthy skin	0.69	0.68

Pz: ratio of colony diameter to total diameter of the colonies plus the precipitation zone; PV: pityriasis versicolor; SD: seborrheic dermatitis.

<sup>a</sup> Not statistically significant difference ( $P=.1$ ).

strains were isolated from individuals without any dermatological disease and showed a mean Pz value the same as that of the healthy individuals included in our research. Table 4 shows the phospholipase activity of *M. sympodialis* strains isolates from patients with PV or SD, from sites with and without evident skin lesions.

After 15 days of incubation, the strains isolated from disease (PV and SD patients), both from healthy skin and lesioned skin, had the same Pz value, while, after 20 days, the strains isolated from the lesioned skin of patients with SD showed higher phospholipasic activity than the strains isolated from healthy skin (but not significantly  $P=0.1$ ) (Table 4).

## Discussion

Extracellular phospholipases are considered virulence factors for many pathogenic bacteria and protozoa such as *Clostridium* species, *Listeria monocytogenes*, *Pseudomonas* species, *Staphylococcus aureus*, *Mycobacterium tuberculosis*, *Toxoplasma gondii* and *Entamoeba histolytica* and for some fungi such as *Candida albicans*, *Aspergillus fumigatus* or *Cryptococcus neoformans*.<sup>10</sup>

Barret-Bee et al.<sup>1</sup> in particular, were the first to correlate the production of phospholipase by *C. albicans* with its pathogenic nature, demonstrating that isolates with a high pathogenic potential (high level of adhesion to oral epithelial cells and greater pathogenicity for the mouse) had higher phospholipasic activity than yeasts with a low pathogenic potential. In addition, *C. albicans* blood isolates have shown greater *in vitro* phospholipasic activity than oral isolates from healthy patients.<sup>16</sup> Similarly, it has been shown that isolates of *C. neoformans* taken from infections in AIDS patients show a higher level of phospholipasic activity than isolates taken from non AIDS patients or from bird droppings.<sup>33,34</sup>

For the *Malassezia* genus some authors have observed the production of different extracellular enzymes such as lipase and phospholipase.<sup>7,17,18,24</sup>

Using the Price method, extra-cellular phospholipasic activity was shown *in vitro* on strains of *M. furfur* isolated from human skin lesions.<sup>22,27</sup> In addition, higher phospholipasic activity was observed in isolates of *M. pachydermatis* from skin lesions in dogs than in those from healthy skin.<sup>5,6</sup>

The method used in this study to highlight the production of phospholipase by yeasts belonging to the *Malassezia* genus proved simple to apply and produced valid results for the species *M. furfur*,

*M. pachydermatis* and *M. sympodialis*, while the strains belonging to the species *M. obtusa*, *M. slooffiae*, *M. globosa*, *M. restricta*, had difficulty developing in the chosen medium so that it was not possible to measure phospholipase activity. To measure the phospholipase activity of *Malassezia*, ten days at least of incubation at 32 °C were needed, unlike the two or six days needed for *C. albicans* and *C. neoformans*, respectively.<sup>28,34</sup>

These results confirm those of other authors<sup>7</sup> and indicate that the yeasts of the *Malassezia* genus produce phospholipase more slowly than *C. albicans* and *C. neoformans*.

For *M. sympodialis* it was possible to compare the production of phospholipase (Pz value) with the origin of the strains. Isolates of *M. sympodialis* taken from patients with PV showed significantly higher phospholipase activity ( $P=0.04$ ) than those isolated from healthy individuals. Strains isolated from patients with SD and from healthy individuals did not show significant differences ( $P=0.08$ ) in the production of phospholipase. Strains isolated from the same patient, from healthy skin and from lesions, had the same Pz on average, both in the case of pityriasis versicolor and of seborrheic dermatitis. It is probable that the same strain colonises the entire individual but producing lesions is only able in some parts of the body, probably in relation to specific local conditions connected with the production of cutaneous lipids and/or alterations of the skin "ecosystem".

These results are only partially comparable with previous works since only a limited amount of data in the literature refers to new species of *Malassezia*.

However, because egg-yolk contains substrates for both phospholipases (phospholipids) and lipases (triglycerides), the egg-yolk-based assay is not entirely specific (although the Pz value correlates with hydrolysis of phosphatidylcholine<sup>26</sup>) and should be considered as an initial screening which requires further confirmation using more specific enquiry such as radiometric or colorimetric methods.

This study suggests however that the phospholipase activity of *Malassezia* may play a role in the onset of skin lesions, especially in the case of PV, even though phospholipases should be considered as only one of the many factors involved in the complex interaction between the yeast and its host leading to the development of skin lesions. More in-depth studies will be needed to understand the pathogenic role played by the enzymes secreted by *Malassezia* spp.

## Conflict of interest

The authors report no conflict of interest.

## References

- Barret-Bee K, Hayes KY, Wilson RG, Ryley JF. A comparison of phospholipase activity, cellular adherence and pathogenicity of yeasts. *J Gen Microbiol.* 1985;131:1217–22.
- Birch M, Robson G, Law D, Denning DW. Evidence of multiple extracellular phospholipase activities in *Aspergillus fumigatus*. *Infect Immun.* 1996;64:751–5.
- Cabañes FJ, Theelen B, Castellá G, Boekhout T. Two new lipid-dependent *Malassezia* species from domestic animals. *FEMS Yeast Res.* 2007;7:1064–76.
- Cabañes FJ, Vega S, Castellá G. *Malassezia cuniculi* sp. nov., a novel yeast species isolated from rabbit skin. *Med Mycol.* 2011;49:40–8.
- Cafarchia C, Gasser RB, Latrofa MS, Parisi A, Campbell BE, Otranto D. Genetic variants of *Malassezia pachydermatis* from canine skin: body distribution and phospholipase activity. *FEMS Yeast Res.* 2008;8:451–9.
- Cafarchia C, Otranto D. Association between phospholipase production by *Malassezia pachydermatis* and skin lesions. *J Clin Microbiol.* 2004;42:4868–9.
- Coutinho SD, Paula CR. Proteinase, phospholipase, hyaluronidase and chondroitin-sulphatase production by *Malassezia pachydermatis*. *Med Mycol.* 2000;38:73–6.
- Crespo Erchiga V, Ojeda A, Vera A, Crespo A, Sánchez F, Guého E. Mycology of pityriasis versicolor. *J Mycol Méd.* 1999;9:143–8.
- Crespo Erchiga V, Ojeda Martos A, Vera Casañó A, Crespo Erchiga A, Sanchez Fajardo F. *Malassezia globosa* as the causative agent of pityriasis versicolor. *Br J Dermatol.* 2000;143:799–803.
- Ghannoum MA. Potential role of phospholipases in virulence and fungal pathogenesis. *Clin Microbiol Rev.* 2000;13:122–43.
- Guého E, Midgley G, Guillot J. The genus *Malassezia* with description of four new species. *Antonie van Leeuwenhoek.* 1996;69:337–55.
- Guillot J, Guého E, Lesourd M, Midgley G, Chevrier G, Dupont B. Identification of *Malassezia* species: a practical approach. *J Mycol Méd.* 1996;6:103–10.
- Guillot J, Guého E. The diversity of *Malassezia* yeasts confirmed by rRNA sequence and nuclear DNA comparisons. *Antonie Van Leeuwenhoek.* 1995;67:297–314.
- Hernández F, Méndez LJ, Bazán E, Arévalo A, Valera A, López R. Especies de *Malassezia* asociadas a diversas dermatosis y a piel sana en población mexicana. *Rev Iberoam Micol.* 2003;20:141–4.
- Hirai A, Kano R, Makimura K, Duarte ER, Hamdan JS, Lachance MA, et al. *Malassezia nana* sp. nov., a novel lipid-dependent yeast species isolated from animals. *Int J Syst Evol Microbiol.* 2004;54:623–7.
- Ibrahim AS, Mirbod F, Filler SG, Banno Y, Cole GT, Kitajima Y, et al. Evidence implicating phospholipase as a virulence factor of *Candida albicans*. *Infect Immun.* 1995;63:1993–8.
- Juntachai W, Oura T, Murayama SY, Kajiwara S. The lipolytic enzymes activities of *Malassezia* species. *Med Mycol.* 2009;47:477–84.
- Mancianti F, Rum A, Nardoni S, Corazza M. Extracellular enzymatic activity of *Malassezia* spp. isolates. *Mycopathologia.* 2001;149:131–5.
- Mayser P, Haze P, Papavassilis C, Pickel M, Gruender K, Guého E. Differentiation of *Malassezia* species: selectivity of Cremophor EL, castor oil and ricinoleic acid for *M. furfur*. *Br J Dermatol.* 1997;137:209–13.
- Mayser P, Laabs S, Heuer KU, Gründer K. Detection of extracellular phospholipase activity in *Candida albicans* and *Rhodotorula rubra*. *Mycopathologia.* 1996;135:149–55.
- Mayser P, Wille G, Imkamp A, Thoma W, Arnold N, Monsees T. Synthesis of fluorochromes and pigments in *Malassezia furfur* by use of tryptophan as the single nitrogen source. *Mycoses.* 1998;41:265–71.
- Muhsin TM, Aubaid AH, Al-Duboom AH. Extracellular enzyme activities of dermatophytes and yeast isolates on solid media. *Mycoses.* 1997;40:465–9.
- Nieverth M, Korting HC. Phospholipases of *Candida albicans*. *Mycoses.* 2001;44:361–7.
- Plotkin LL, Squiquera L, Mathov I, Galimberti R, Leoni J. Characterization of the lipase activity of *Malassezia furfur*. *J Med Vet Mycol.* 1996;34:43–8.
- Polak A. Virulence of *Candida albicans* mutants. *Mycoses.* 1992;35:9–16.
- Price MF, Wilkinson ID, Gentry LO. Plate method for detection of phospholipase activity in *Candida albicans*. *Sabouraudia.* 1982;20:7–14.
- Riciputo RM, Oliveri S, Micali G, Sapuppo A. Phospholipase activity in *Malassezia furfur* pathogenic strains. *Mycoses.* 1996;39:233–5.
- Samaranayake LP, Reaside JM, MacFarlane TW. Factors affecting the phospholipase activity of *Candida* species in vitro. *Sabouraudia.* 1984;22:201–7.
- Sugita T, Tajima M, Takashima M, Amaya M, Saito M, Tsuboi R, et al. A new yeast, *Malassezia yamatoensis*, isolated from a patient with seborrheic dermatitis, and its distribution in patients and healthy subjects. *Microbiol Immunol.* 2004;48:579–83.
- Sugita T, Takashima M, Kodama M, Tsuboi R, Nishikawa A. Description of a new yeast species, *Malassezia japonica*, and its detection in patients with atopic dermatitis and healthy subjects. *J Clin Microbiol.* 2003;41:4695–9.
- Sugita T, Takashima M, Shinoda T, Suto H, Unno T, Tsuboi R, et al. New yeast species, *Malassezia dermatis*, isolated from patients with atopic dermatitis. *J Clin Microbiol.* 2002;40:1363–7.
- Tragiannidis A, Bisping G, Koehler G, Groll AH, Minireview. *Malassezia* infections in immunocompromised patients. *Mycoses.* 2010;53:187–95.
- Vidotto V, Leone R, Sinicco A, Ito-kuwa S, Criseo G. Comparison of phospholipase production in *Cryptococcus neoformans* isolates from AIDS patients and bird droppings. *Mycopathologia.* 1998;142:71–6.
- Vidotto V, Sinicco A, Di Fraia D, Cardaropoli S, Aoki S, Ito-Kuwa S. Phospholipase activity in *Cryptococcus neoformans*. *Mycopathologia.* 1996;136:119–23.
- Xu J, Saunders CW, Hu P, Grant RA, Boekhout T, Kuramae EE, et al. Dandruff-associated *Malassezia* genomes reveal convergent and divergent virulence traits shared with plant and human fungal pathogens. *Proc Natl Acad Sci U S A.* 2007;104:18730–5.