



Screening for ligninolytic enzymes in autochthonous fungal strains from Argentina isolated from different substrata

Mario Carlos Nazareno Saparrat¹, María Jesús Martínez²,
Marta Noemí Cabello¹ & Angélica Margarita Arambarri¹

¹Instituto de Botánica Spegazzini, Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, Argentina; ²Centro de Investigaciones Biológicas, Consejo Superior de Investigaciones Científicas, Madrid, Spain

Summary

The production of different extracellular ligninolytic enzymes was studied in autochthonous fungal strains from Argentina isolated from litter derived from hydrocarbon-polluted sites and from basidiocarps growing on wood in forests. The strains tested were cultivated in a carbon-limited medium with shaking. Laccase activity reached higher levels than aryl-alcohol oxidase and manganese-dependent peroxidase activities in liquid cultures from different fungi. No lignin peroxidase activity was found in any strain assayed. Some species are reported for the first time as producers of different ligninolytic enzymes.

Key words

Aryl-alcohol oxidase, Basidiomycetes, Deuteromycetes, Laccase, Lignin peroxidase, Manganese-peroxidase

Enzimas ligninolíticas de cepas fúngicas autóctonas de Argentina aisladas de diferentes sustratos

Resumen

Se estudió la producción de diferentes enzimas ligninolíticas extracelulares en cepas fúngicas autóctonas de Argentina aisladas a partir de materia orgánica de sitios contaminados con hidrocarburos y basidiocarpos desarrollados sobre restos leñosos de bosques. Las diferentes cepas estudiadas se cultivaron en un medio limitado en carbono bajo agitación. Se detectaron niveles superiores de actividad lacasa en relación a los correspondientes para las actividades extracelulares de aril-alcohol oxidasa y peroxidasa dependiente de manganeso en los cultivos líquidos de diferentes hongos. No se detectó actividad lignina peroxidasa en ninguno de los aislamientos analizados. Diferentes especies son citadas por primera vez como productoras de diferentes enzimas ligninolíticas. *Palabras clave:* Aril-alcohol oxidasa, basidiomycetes, deuteromycetes, lacasa, lignina peroxidasa, peroxidasa dependiente de manganeso.

Palabras clave

Aril-alcohol oxidasa, Basidiomycetes, Deuteromycetes, Lacasa, Lignina peroxidasa, Peroxidasa dependiente de manganeso

Lignin is a recalcitrant heteropolymer of phenylpropanoid units present in woody plant tissues, that confers them rigidity and resistance to biological attack [1]. In order to depolymerize and mineralize lignin, white-rot fungi have developed an oxidative and unspecific system including extracellular enzymes, low molecular weight metabolites and activated oxygen species [2-5]. Due to the lack of specificity of the system involved in the lignin depolymerization, white-rot basidiomycetes and their enzymes are being studied for their application on the degradation of aromatic pollutants causing environmental problems [6-9].

The extracellular enzymatic systems include ligninolytic peroxidases, laccases and oxidases responsible for the production of extracellular hydrogen peroxide (H₂O₂) [10]. Those enzyme systems exhibit differential characteristics depending on the species, strains and culture conditions [11-13]. Among peroxidases, lignin peroxidase (LiP) is able to oxidize directly non-phenolic units whereas manganese peroxidase (MnP) and laccase oxidize prefe-

Dirección para correspondencia:

Dr. Mario Carlos Nazareno Saparrat
Instituto de Botánica Spegazzini
Facultad de Ciencias Naturales y Museo
Universidad Nacional de La Plata, 53 # 477
1900-La Plata, Argentina
Tel./fax: +54 221 4219845
E-mail: speg_lps@museo.fcnym.unlp.edu.ar

Aceptado para publicación el 23 de Mayo de 2002

rentially phenolic units, but also act on non-phenolic units in the presence of mediators [14,15]. Recently there have been reports of a new peroxidase type in *Pleurotus* and *Bjerkandera* species which shares catalytic properties with LiP and MnP [16-17] and is also able to oxidize azo-dyes not oxidized by MnP or LiP [18]. On the other hand, two extracellular oxidases, glyoxal and aryl-alcohol (AAO) oxidases, have been reported for extracellular H₂O₂ production [19,20].

Several screening works about ligninolytic enzymes have been carried out mainly in white-rot basidiomycetes [21-24]. However, other fungi, representatives of different taxonomic and ecophysiological groups, are able to degrade lignocellulosic substrata, mineralize ¹⁴C-milled wood lignin [25] and produce ligninolytic enzymes [13,26].

The aim of the present work was to produce different extracellular enzymes involved in lignin degradation and detoxification of aromatic pollutants and to determine the ability of several argentinian autochthonous basidiomycetes and deuteromycetes strains, isolated from organic matter of hydrocarbon-polluted sites and basidiomycetes developed on wood, to produce these enzymes.

Materials and methods

Fungal strains. The strains used for this work were isolated from different sources/substrata and habitats from Argentina (Table 1). They belong to the culture collection of Spegazzini Institute (CLPS). Stock cultures of the basidiomycetes strains were kept at 4 °C on 2% (w/v) malt-agar slants supplemented with yeast extract (0.4%) and *Populus* spp. wood chips. Cultures of the deuteromycetes were maintained on agar slants (Table 1) at 4 °C.

Medium and culture conditions. The production of extracellular ligninolytic enzymes was carried out in the

modified Czapek Dox medium [23]. Homogenized mycelium from 5-10 day-old cultures was used to inoculate 250 ml Erlenmeyer flasks containing 50 ml of medium (3-5 mg of dry weight/ml of inoculum). Four replicate flasks were incubated at 25 °C in a rotary shaker at 150 rev/min.

Analytical methods. The cultures were harvested at the 7th and 21st day of incubation. Each sample was centrifuged (10,000 x g for 30 min) at 4 °C. The mycelial pellet was dried at 60 °C and weighed to estimate the fungal biomass. The supernatant of the liquid culture was kept for enzyme assays.

Enzyme assays. AAO (EC 1.1.3.7) activity was estimated by the veratraldehyde formation from 5 mM veratryl alcohol (VA) (Fluka) in 100 mM phosphate buffer at pH 6.0 (ϵ_{310} : 9,300/Mcm) [20]. Laccase (EC 1.10.3.2) activity was measured with 5 mM 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid) (ABTS) (Sigma) in 100 mM acetate buffer at pH 5.5 (ϵ_{436} of the ABTS cation radical: 29,300/Mcm) [23]. LiP (EC 1.11.1.14) activity was determined by H₂O₂-dependent veratraldehyde formation from 2 mM VA in 100 mM sodium tartrate buffer at pH 3.0, with H₂O₂ 0.4 mM [27]. MnP (EC 1.11.1.13) activity was estimated using as substrate 0.01 % phenol-red (Sigma) in 100 mM sodium tartrate, pH 5.0 [28], in presence of 0.1 mM H₂O₂ and 1 mM MnSO₄.

The enzyme reactions were carried out in duplicate and controls were performed without the addition of the enzyme or H₂O₂ for oxidase and peroxidase assays, respectively. Based on preliminary studies, optimal assay times were known to fall in the linear range of enzyme kinetics. All oxidation rates were determined at 25 °C using a Beckman DU 640 u.v.-visible spectrophotometer. One enzymatic activity unit (U) was defined as the amount of enzyme that transforms 1 μ mol of substrate/min.

Table 1. Fungal strains used in this study.

Fungal species and taxonomic groups	CLPS ^a	Substrate/habitat and collection site
Basidiomycetes		
<i>Amauroderma boleticeum</i> (Pat. And Gaill.) Torr.	157	Decaying wood of subtropical rain forests ^{Ga}
<i>Auricularia</i> sp.	550	Decaying wood of subtropical rain forests ^A
<i>Corioloopsis rigida</i> (Berk. Et Mont.) Murrill	232	Rotten wood of subtropical rain forests ^{Ga}
<i>Cyathus striatus</i> (Hudson) Hoffm.	381	Rotten wood of subtropical rain forests ^{Ga}
<i>Grammothele subargentea</i> (Speg.) Rajch.	436	Trunk of living tree <i>Angiosperm</i> of subtropical rain forests ^{Ga}
<i>Loweporus lividus</i> (Kalchbrenner) Wright	289	Rotten wood of subtropical rain forests ^{Ga}
<i>Peniophora albobadia</i> (Schw.: Fr.) Boidin	285	Decaying wood of subtropical rain forests ^{Ga}
<i>Phanerochaete septocystidia</i> (Burt.) Erikss	288	Rotten wood of subtropical rain forests ^{Ga}
<i>Phanerochaete tuberculata</i> (Karst.) Parm.	435	Rotten wood of subtropical rain forests ^{Ga}
<i>Phellinus gilvus</i> (Schw.) Pat. var. <i>licnoides</i> (Mont.)	156	Rotten wood of subtropical rain forests ^{Ga}
<i>Phellinus linteus</i> (Berk. Et Curt.) Teng.	338	Trunk of living tree of subtropical rain forests ^{Ga}
<i>Pleurotus laciniatocrenatus</i> (Speg.) Speg.	39	Trunk of living tree <i>Taxodium</i> sp. of urban forest area ^{Pe}
<i>Pycnoporus sanguineus</i> (L.: Fr.) Murr.	163	Trunk of living tree <i>Leguminosae</i> of subtropical rain forests ^{Ga}
<i>Trametes pavonia</i> (Hook.) Ryv.	437	Rotten trunk of subtropical rain forests ^{Ga}
<i>Trametes subectypus</i> (Murr.) Gibn.	342	Rotten wood of subtropical rain forests ^{Ga}
<i>Trametes villosa</i> (Fr.) Kreisel	233	Rotten trunk of subtropical rain forests ^{Ga}
Deuteromycetes		
<i>Alternaria alternata</i> (Fries: Fries) Keissler	267 ^b	Floating litter collected in hydrocarbon-polluted water ^S
<i>Beltrania rhombica</i> Penz	272 ^c	Floating litter collected in hydrocarbon-polluted water ^S
<i>Cladosporium cladosporioides</i> (Fresen.) de Vries	556 ^d	Hydrocarbon-polluted soil ^F
<i>Epicoccum purpurascens</i> Ehrenb. Ex Schlecht.	554 ^d	Hydrocarbon-polluted soil ^F
<i>Fusarium solani</i> (Martius) Saccardo	493 ^c	Hydrocarbon-polluted soil ^F
<i>Graphium putredinis</i> (Corda) Hughes	423 ^c	Hydrocarbon-polluted soil ^F
<i>Minimidochium parvum</i> Cabello, Arambarri and Cazau	548 ^d	Floating litter collected in hydrocarbon-polluted water ^S
<i>Penicillium chrysogenum</i> Thom	495 ^c	Crude oil ^F
<i>Phaeoisaria clematidis</i> (Fuckel) Hughes	154 ^d	Floating litter collected in hydrocarbon-polluted water ^S
<i>Tetraploa aristata</i> Berkeley et Broome	419 ^d	Hydrocarbon-polluted organic matter floating in freshwater ^S
<i>Trichoderma saturnisporum</i> Hammill	264 ^d	Floating litter collected in hydrocarbon-polluted water ^S

^aCLPS strain no. Culture media deuteromycetes type: ^bcorn-meal agar medium; ^cczapek with 1% (v/v) crude oil; ^d2% (w/v) malt-agar; ^epotato extract-agar. Collection sources: ^AAristóvalo del Valle (Province of Misiones), Argentina; ^EEnsenada (Province of Buenos Aires), Argentina; ^{Ga}Garupá (Province of Misiones), Argentina; ^{Pe}Pereyra Park (Province of Buenos Aires), Argentina; ^SSantiago river (Province of Buenos Aires), Argentina; ^{SA}Santa Ana (Province of Misiones), Argentina.

Results and discussion

The ability of different argentinian fungal strains belonging to basidiomycetes and deuteromycetes to produce extracellular ligninolytic enzymes in the modified Czapek Dox medium under shaking was screened. All the fungal strains studied produced abundant growth under the culture conditions assayed with a yield between 200 and 600 mg/100 ml of medium. The highest biomass values were obtained with the cultures of *Cyathus striatus* - at 21 days of incubation -. As a consequence of the utilization of a C-limited medium, the biomass obtained at 7 days (trophophase) was higher than at 21 days (idiophase).

No ligninolytic enzymatic activities were detected in the basidiomycetes *Cyathus striatus*, *Loweporus lividus*, and *Phellinus linteus* and the deuteromycetes *Alternaria alternata*, *Beltrania rhombica*, *Cladosporium cladosporioides*, *Epicoccum purpurascens*, *Penicillium chrysogenum* and *Trichoderma saturnisporum* in the culture conditions assayed.

LiP activity was not detected in any of the strains studied. Previous screening, including the basidiomycete *Phanerochaete chrysosporium*, reported similar results in shake or static conditions [21,23]. It is known that *P. chrysosporium* needs specific growth conditions to produce LiP, including nitrogen limited medium and O₂ saturated atmosphere [27]. It is possible that the culture conditions used in this study could affect LiP expression although other fungi produce LiP in air-atmosphere and non-limited nitrogen medium [29]. In this sense, it is possible that the absence of VA as LiP inducer in the culture medium, or the inhibition of this activity by aromatic compounds present in the culture, are responsible for the negative results [30-31].

The ligninolytic enzymatic activities, AAO, laccase and MnP, detected after 7 and 21 days of incubation in the different strains studied are shown in figures 1, 2 and 3, respectively. Note that the activity levels obtained did not necessarily reflect the optimum production for each strain.

The AAO activity was observed only in the 25 % of the strains tested (Figure 1), reaching the highest levels - more than 5 mU/ml - in *Amauroderma boleticeum* at 7 days and in *Pleurotus laciniatocrenatus* at 21 days of incubation. Extracellular AAO has been characterized from the culture liquid of *Pleurotus* and *Bjerkandera* species [20,32-35] and from the mycelial extract of *Phanerochaete chrysosporium* [36]. The AAO catalyzes the oxidation of a broad number of aromatic alcohols, chemically related to lignin, to aldehydes [20] and it participates in the continuous H₂O₂ production through a system based on aromatic aldehydes redox cycling [37-38].

Extracellular laccase activity was detected in most of the basidiomycetes strains studied (Figure 2); *Grammothele subargentea* reached the highest activity (170 mU/ml). According to the results obtained in previous screening works [21,23], laccase was the main ligninolytic activity detected. This activity was also produced by some of the deuteromycetes strains tested (*Minimidochium parvum* and *Tetraploa aristata*). Laccase is a copper-containing oxidase involved in lignin biodegradation and secreted by most basidiomycetes [10,21,23]. However, this enzyme also participates in other physiological processes, as conidial pigmentation, morphogenesis, pathogenesis and detoxification of phenolic compounds, and it has also been characterized from different ascomycetes and deuteromycetes [39]. Recently, the participation of this enzyme in oxygen activation during the oxidation of hydroquinones [40] and its syner-




Figure 1. AAO activities in the culture fluid of different fungi grown in the modified Czapek Dox medium (mean values).




Figure 2. Laccase activities in the culture fluid of different fungi grown in the modified Czapek Dox medium (mean values).




Figure 3. MnP activities in the culture fluid of different fungi grown in the modified Czapek Dox medium (mean values).

gistic action with AAO to produce hydroxyl radical [41], a low molecular weight compound involved in the initial attack to lignocellulose when ligninolytic enzymes can not penetrate plant cell walls [42], has been reported. On the other hand, the addition of aromatic compounds to the fungal cultures may induce the expression of different laccase isoforms [43,44]. In this work, the ligninolytic enzymatic activities have been studied in a carbon-limited medium, in absence of inducers, suggesting that the ligninolytic enzymes reported here could be constitutive.

Laccase production, induced by xyloidine, has been already reported in *Pycnoporus sanguineus* [45]. However, it is necessary to mention that some fungi, as *Cladosporium cladosporioides* and *Penicillium chrysogenum*, only secrete laccase in presence of inducers such as lignin derived compounds, humic acids or xyloidine [46-47].

As laccase, MnP production was detected in numerous strains studied (Figure 3). However, the MnP levels were comparatively lower. These results agree with those reported from other fungi using phenol red as substrate [23,26]. This study is the first report on the production of peroxidases related to ligninolytic systems by fungal species belonging to different taxonomic and ecophysiological groups. Except *Auricularia* sp. [48], *Fusarium solani* [26] and *Pycnoporus sanguineus* [49], the fungi listed in figure 3 are reported for the first time as producers of extracellular MnP. This enzyme could play an important role in lignin biodegradation in fungi lacking LiP activity. *Pleurotus* species can mineralize lignin from ¹⁴C-wheat straw and a manganese oxidizing peroxidase is involved in the process [50].

Most basidiomycetes with ligninolytic enzymatic activity secrete laccase and MnP simultaneously. However, the production of AAO, laccase and MnP has been detected only in *Auricularia* sp., *Grammothele subargentea*, *Phellinus gilvus* and *Pleurotus laciniatocrenatus*. In deuteromycetes with enzymatic activity, MnP has been the enzyme found in most species, although laccase and AAO are also secreted by some fungi.

Basidiomycetes, which produce ligninolytic enzymes, are associated with wood rots and lignin degradation [10]. However, the presence of laccase and peroxidase activities in *Minimidochium parvum* and *Tetraploa aristata*, deuteromycetes isolated from soil and floating litter in water polluted with crude-oil from Argentina, could be involved in the degradation of xenobiotics and detoxification of polluted systems since both enzymes are involved in the biodegradation of recalcitrant aromatic compounds [6-7,51].

The authors wish to thank Lic. Bucsinzky, A.M.M. (CONICET technician) who is in charge of the culture collection at the Spegazzini Institute, and Lic. Tournier, H.A. (Fac. Ciencias Médicas UNLP) for their excellent technical assistance. This chapter has benefited from the constructive criticism of anonymous reviewers. Saparrat, M.C.N. is a recipient of a scholarship from CONICET. Cabello, M.N. is a researcher from CIC. Arambarri, A.M. is a researcher from CONICET. This research was supported by Grants from CONICET and CIC, Argentina.

References

- Higuchi T. Lignin biochemistry: biosynthesis and biodegradation. Wood Sci Technol 1990; 24: 23-63.
- Schoemaker HE. On the chemistry of lignin degradation. Recueil des Travaux Chimiques des Pays-Bas 1990; 109: 255-272.
- Shimada M, Higuchi T. Microbial, enzymatic and biomimetic degradation of lignin. In: Hon DNS, Shiraiishi N (Eds.) Wood and cellulosic chemistry. New York, Marcel Dekker, 1991: 557-619.
- de Jong E, Field JA, Dings JAFM, Wijnberg JBPA, de Bont JAM. De novo biosynthesis of chlorinated aromatics by the white-rot fungus *Bjerkandera* sp BOS55. Formation of 3-chloro-anisaldehyde from glucose. FEBS Lett 1992; 305: 220-224.
- Guillén F, Martínez MJ, Muñoz C, Martínez AT. Quinone redox cycling in the ligninolytic fungus *Pleurotus eryngii* leading to extracellular production of superoxide anion radical. Arch Biochem Biophys 1997; 339: 190-199.
- Hammel KE. Oxidation of aromatic pollutants by lignin-degrading fungi and their extracellular peroxidases. In: Sigel H, Sigel A (Eds.) Metal ions in biological systems. Vol. 28. Degradation of environmental pollutants by microorganisms and their metalloenzymes. New York, Marcel Dekker, 1992: 41-60.
- Field JA, de Jong E, Feijoo-Costa G, de Bont JAM. Screening for ligninolytic fungi applicable to the biodegradation of xenobiotics. Trends Biotechnol 1993; 11: 44-49.
- Paszczynski A, Crawford RL. Potential for bioremediation of xenobiotic compounds by the white-rot fungus *Phanerochaete chrysosporium*. Biotechnol Progress 1995; 11: 368-379.
- Novotny C, Vyas BRM, Erbanova P, Kubatova A, Sasek V. Removal of PCBs by various white rot fungi in liquid cultures. Folia Microbiol (Prague) 1997; 42: 136-140.
- Hatakka A. Lignin-modifying enzymes from selected white-rot fungi - Production and role in lignin degradation. FEMS Microbiol Rev 1994; 13: 125-135.
- Rogalski J, Lundell TK, Leonowicz A, Hatakka AI. Influence of aromatic compounds and lignin on production of ligninolytic enzymes by *Phlebia radiata*. Phytochemistry 1991; 30: 2869-2872.
- Hammer E, Schauer F. Fungal hydroxylation of dibenzofuran. Mycol Res 1997; 101: 433-436.
- Heinzkill M, Messner K. The ligninolytic system of fungi. In: Anke T (Ed.) Fungal Biotechnology. Weinheim, Chapman & Hall, 1997: 213-227.
- Hammel KE, Tardone PJ, Moen MA, Price LA. Biomimetic oxidation of nonphenolic lignin models by Mn(III): new observations on the oxidizability of guaiacyl and syringyl substructures. Arch Biochem Biophys 1989; 270: 404-409.
- Bourbonnais R, Paice MG. Oxidation of non-phenolic substrates. An expanded role for laccase in lignin biodegradation. FEBS Lett 1990; 267: 99-102.
- Martínez MJ, Ruiz-Dueñas FJ, Guillén F, Martínez AT. Purification and catalytic properties of two manganese-peroxidase isoenzymes from *Pleurotus eryngii*. Eur J Biochem 1996; 237: 424-432.
- Mester T, Field JA. Characterization of a novel manganese peroxidase-lignin peroxidase hybrid isozyme produced by *Bjerkandera* species strain BOS55 in the absence of manganese. J Biol Chem 1998; 273: 15412-15417.
- Heinfling A, Ruiz-Dueñas FJ, Martínez MJ, Bergbauer M, Szewzyk U, Martínez AT. A study on reducing substrates of manganese-oxidizing peroxidases from *Pleurotus eryngii* and *Bjerkandera adusta*. FEBS Lett 1998; 428: 141-146.
- Kersten PJ. Glyoxal oxidase of *Phanerochaete chrysosporium*: its characterization and activation by lignin peroxidase. Proc Natl Acad Sci USA 1990; 87: 2936-2940.
- Guillén F, Martínez AT, Martínez MJ. Substrate specificity and properties of the aryl-alcohol oxidase from the ligninolytic fungus *Pleurotus eryngii*. Eur J Biochem 1992; 209: 603-611.
- Szklarz GD, Antibus RK, Sinsabaugh RL, Linkins AE. Production of phenol oxidases and peroxidases by wood-rotting fungi. Mycologia 1989; 81: 234-240.
- de Jong E, de Vries FP, Field JA, van der Zwan RP, de Bont JAM. Isolation and screening of basidiomycetes with high peroxidative activity. Mycol Res 1992; 96: 1098-1104.
- Peláez F, Martínez MJ, Martínez AT. Screening of 68 species of basidiomycetes for enzymes involved in lignin degradation. Mycol Res 1995; 99: 37-42.
- Heinzkill M, Bech L, Halkier T, Schneider P, Anke T. Characterization of laccases and peroxidases from wood-rotting fungi (Family *Coprinaceae*). Appl Environ Microbiol 1999; 64: 1601-1606.
- Rodríguez A, Perestelo F, Carnicero A, et al. Degradation of natural lignins and lignocellulosic substrates by soil-inhabiting fungi imperfecti. FEMS Microbiol Ecol 1996; 21: 213-219.
- Saparrat MCN, Martínez MJ, Tournier HA, Cabello MN, Arambarri AM. Production of ligninolytic enzymes by *Fusarium solani* strains isolated from different substrata. World J Microbiol Biotechnol 2000; 16: 799-803.
- Tien M, Kirk TK. Lignin peroxidase of *Phanerochaete chrysosporium*. Methods Enzymol 1988; 161: 238-248.
- Michel FC, Dass SB, Grulke EA, Reddy CA. Role of manganese peroxidases and lignin peroxidases of *Phanerochaete chrysosporium* in the decolorization of kraft bleach plant effluent. Appl Environ Microbiol 1991; 57: 2368-2375.
- Kaal EEJ, de Jong E, Field JA. Stimulation of ligninolytic peroxidase activity by nitrogen nutrients in the white rot fungus *Bjerkandera* sp strain BOS55. Appl Environ Microbiol 1993; 59: 4031-4036.

30. Buswell JA, Odier E. Lignin biodegradation. *Crit Rev Biotechnol* 1987; 6: 1-60.
31. Vares T, Lundell TK, Hatakka AI. Novel heme-containing enzyme possibly involved in lignin degradation by the white-rot fungus *Junghuhnia separabilima*. *FEMS Microbiol Lett* 1992; 99: 53-58.
32. Bourbonnais R, Paice MG. Veratryl alcohol oxidases from the lignin degrading basidiomycete *Pleurotus sajor-caju*. *Biochem J* 1988; 255: 445-450.
33. Muheim A, Waldner R, Leisola MSA, Fiechter A. An extracellular aryl-alcohol oxidase from the white-rot fungus *Bjerkandera adusta*. *Enzyme Microb Technol* 1990; 12: 204-209.
34. Sannia G, Limongi P, Cocca E, Buonocore F, Nitti G, Giardina P. Purification and characterization of a veratryl alcohol oxidase enzyme from the lignin degrading basidiomycete *Pleurotus ostreatus*. *Biochim Biophys Acta* 1991; 1073: 114-119.
35. Varela E, Martínez AT, Martínez MJ. Molecular cloning of aryl-alcohol oxidase from *Pleurotus eryngii*, an enzyme involved in lignin degradation. *Biochem J* 1999; 341: 113-117.
36. Asada Y, Watanabe A, Ohtsu Y, Kuwahara M. Purification and characterization of an aryl-alcohol oxidase from the lignin-degrading basidiomycete *Phanerochaete chrysosporium*. *Biosci Biotechnol Biochem* 1995; 59: 1339-1341.
37. Guillén F, Evans CS. Anisaldehyde and veratraldehyde acting as redox cycling agents for H₂O₂ production by *Pleurotus eryngii*. *Appl Environ Microbiol* 1994; 60: 2811-2817.
38. Guillén F, Martínez AT, Martínez MJ, Evans CS. Hydrogen peroxide-producing system of *Pleurotus eryngii* involving the extracellular enzyme aryl-alcohol oxidase. *Appl Microbiol Biotechnol* 1994; 41: 465-470.
39. Thurston CF. The structure and function of fungal laccases. *Microbiology-UK* 1994; 140: 19-26.
40. Guillén F, Muñoz C, Gómez-Toribio V, Martínez AT, Martínez MJ. Oxygen activation during the oxidation of methoxyhydroquinones by laccase from *Pleurotus eryngii*. *Appl Environ Microbiol* 2000; 66: 170-175.
41. Guillén F, Gómez-Toribio V, Muñoz C, Martínez MJ, Martínez AT. Production of hydroxyl radical by the synergistic action of fungal laccase and aryl alcohol oxidase. *Arch Biochem Biophys* 2000; 382: 142-147.
42. Evans CS, Dutton MV, Guillén F, Veness RG. Enzymes and small molecular mass agents involved with lignocellulose degradation. *FEMS Microbiol Rev* 1994; 13: 235-240.
43. Bollag JM, Leonowicz A. Comparative studies of extracellular fungal laccases. *Appl Environ Microbiol* 1984; 48: 849-854.
44. Muñoz C, Martínez AT, Martínez MJ. Induction and partial purification of laccase from *Pleurotus eryngii*. In: Duarte JC, Ferreira MC, Ander P (Eds.) *Proceedings of the FEMS Symposium on Lignin Biodegradation and Transformation*, Lisbon, 17-21 April. Lisbon, Forbitech, 1993: 173-174.
45. Pointing SB, Jones EBG, Vrijmoed LLP. Optimization of laccase production by *Pycnoporus sanguineus* in submerged liquid culture. *Mycologia* 2000; 92: 139-144.
46. Rodríguez A, Falcón MA, Carnicero A, Perestelo F, de la Fuente G, Trojanowski J. Laccase activities of *Penicillium chrysogenum* in relation to lignin degradation. *Appl Microbiol Biotechnol* 1996; 45: 399-403.
47. Claus H, Filip Z. Degradation and transformation of aquatic humic substances by laccase-producing fungi *Cladosporium cladosporioides* and *Polyporus versicolor*. *Acta Hydrochimica et Hydrobiologica* 1998; 26: 180-185.
48. Hofrichter M, Fritsche W. Depolymerization of low rank coal by extracellular fungal enzyme systems. 1. Screening for low rank coal-depolymerizing activities. *Appl Microbiol Biotechnol* 1996; 46: 220-225.
49. Esposito E, Innocentini LH, Ferraz A, Canhos VP, Durán N. Phenoloxidases and hydrolases from *Pycnoporus sanguineus* (UEC-2050 strain) - Applications. *J Biotechnol* 1993; 29: 219-228.
50. Camarero S, Böckle B, Martínez MJ, Martínez AT. Manganese-mediated lignin degradation by *Pleurotus pulmonarius*. *Appl Environ Microbiol* 1996; 62: 1070-1072.
51. Johannes C, Majcherzyk A. Natural mediators in the oxidation of polycyclic aromatic hydrocarbons by laccase mediator systems. *Appl Environ Microbiol* 2000; 66: 524-528.